The listing of claims presented below replaces all prior versions and listing of claims in the application.

Listing of claims:

- 1. (Withdrawn) A method for the treatment of an individual having a condition characterized by abnormal myocardial cell Na⁺, K⁺ or Ca²⁺ ion levels, said method comprising administering a therapeutically effective amount of one or more ss; 3 β_3 adrenoceptor agonists to said individual.
- 2. (Withdrawn) The method according to claim 1 wherein the condition is selected from the group consisting of heart failure, and myocardial hypertrophy.
- 3. (Currently amended) A method for the treatment of <u>a human an individual</u> suffering from or susceptible to heart failure or myocardial hypertrophy, said method comprising administering a therapeutically effective amount of one or more β_3 adrenoceptor agonists to said <u>human an individual</u>.
- 4. (Currently amended) The method according to claim 3 wherein the <u>human has</u> individual is an individual having one or more clinical symptoms of heart failure or myocardial hypertrophy.
- 5. (Previously presented) The method according to claim 3 wherein the β_3 adrenoceptor agonist is selected from the group consisting of arylethanolamines, aryloxypropanolamines, and trimetoquinols.
- 6. (Previously presented) The method according to claim 3 wherein the β_3 adrenoceptor agonist is selected from the group consisting of BRL37344, BRL 26830, BRL 26830A, BRL 35113, ZD7114, CGP12177, CGP 12177A, CGP-20712A, CL316243, ICI07114, ICI215001, ICI 201651, BRL35135A, BRL28410, N-5984, (R)-N-[4-[2-[[2-Hydroxy-2- (pyridin-3-yl)ethyl]amino]ethyl]phenyl]- 4- [4-(4-trifluoro-methylphenyl)thiazol-2- yl] benzenesulfonamide (L-796568), (R)-N-[4-[2-[[2-hydroxy-2-(3-pyridinyl)- ethyl]amino]ethyl]phenyl]-1-(4-octylthiazol-2-yl)

- yl)-5-indolinesulfonamide (L-755507), L- 770,644, L-766,892, L-757,793, L-796568, LY-377604, Ro 40-2148, SB-220646, SB- 226552, SB-251023, SB-262552, SR 58306, SR 58375, SR 58339, SR 58611, SR 58611A, SR 59119A, GR-265261-X, AD-9677, and agonists of the series 2-(3-indolyl) alkylamino-1-(3-chlorophenyl)ethanols.
- 7. (Original) The method according to claim 3 wherein the β_3 adrenoceptor agonist is BRL37344.
- 8. (Currently amended) The method according to claim 3 wherein the β_3 -drenoceptor agonist further comprises β_1 antagonist activity and or and/or further comprises $\frac{1}{3}$ antagonist activity.
- 9. (Currently amended) The method according to claim 3 further comprising administering one or more β blockers to said <u>human individual</u>.
- 10. (Original) The method according to claim 9 wherein the β blocker is nadolol.
- 11. (Original) The method according to claim 9 wherein the β blocker is a β_1 and/or β_2 adrenoceptor antagonist.
- 12. (Currently amended) The method according to claim 9 wherein the β blocker is administered to said <u>human</u> individual prior to, simultaneously with or subsequent to administration of the one or more β_3 adrenoceptor agonists.
- 13. (Currently amended) The method according to claim 3 wherein said human is suffering from chronic heart failure and further comprising at least partially stabilizing said human individual prior to administration of said β_3 adrenoceptor agonist.
- 14. (Original) The method according to claim 13 wherein said stabilizing comprises treatment with one or more compounds selected from the group consisting of ACE- inhibitors, aldosterone antagonists and β-adrenoceptor antagonists.

- 15. (Withdrawn) A method for treatment of a condition characterized by abnormally high myocardial cell Na+ ion level, said method comprising administration to an individual having said condition of a therapeutically effective amount of one or more β₃- adrenoceptor agonists.
- 16. (Withdrawn) The method according to claim 15 wherein said condition characterized by abnormally high myocardial cell Na⁺ ion level is selected from the group consisting of heart failure, myocardial hypertrophy, and diabetic cardiomyopathy.
- 17. (Withdrawn) Use of one or more β_3 adrenoceptor agonists for the manufacture of a medicament for treatment of an individual having a condition characterized by abnormal myocardial cell Na⁺, K⁺ or Ca²⁺ ion levels.
- 18. (Withdrawn) One or more β_3 -adrenoceptor agonists for use in the treatment of an individual having a condition characterized by abnormal myocardial cell Na⁺, K⁺ or Ca²⁺ ion levels.
- 19. (Withdrawn) Use of one or more β_3 -adrenoceptor agonists for the manufacture of a medicament for treatment of an individual suffering from or susceptible to heart failure or myocardial hypertrophy.
- 20. (Withdrawn) One or more β_3 -adrenoceptor agonists for use in the treatment of an individual suffering from or susceptible to heart failure or myocardial hypertrophy.
- 21. (Withdrawn) A pharmaceutical composition for use in the treatment of an individual having a condition characterized by abnormal myocardial cell Na⁺, K⁺ or Ca²⁺ ion levels, the composition comprising one or more β_3 adrenoceptor agonists together with one or more pharmaceutically acceptable adjuvants, excipients and/or carriers.
- 22. (Withdrawn) A pharmaceutical composition for use in the treatment of an individual suffering from or susceptible to heart failure or myocardial hypertrophy, the composition comprising one or more β_3 adrenoceptor agonists together with one or more pharmaceutically acceptable adjuvants,

excipients and/or carriers.

- 23. (Withdrawn) A pharmaceutical composition comprising one or more β_3 adrenoceptor agonists and one or more β_1 and/or β_2 adrenoceptor antagonists, together with one or more pharmaceutically acceptable adjuvants, excipients and/or carriers.
- 24. (Withdrawn) A method for the extrusion of Na⁺ from a myocardial cell or cells, the method comprising contacting said cell (s) one or more β_3 adrenoceptor agonist(s).
- 25. (Withdrawn) The method according to claim 24 wherein said method comprises Na,K pump stimulation.